

AD-A186 433

STRONG REPRESENTATION OF WEAK CONVERGENCE(U) NORTH  
CAROLINA UNIV AT CHAPEL HILL DEPT OF STATISTICS  
Z D BAI ET AL JUN 87 TR-186 AFOSR-TR-87-1334

1/1


UNCLASSIFIED

F49628-85-C-8144

F/G 12/3

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A186 433

FILE

2


## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY NA		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; Distribution Unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 87 - 1354	
4. PERFORMING ORGANIZATION REPORT NUMBER Technical Report No. 186		7a. NAME OF MONITORING ORGANIZATION AFOSR/NM	
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina		7b. ADDRESS (City, State and ZIP Code) Building 410 Bolling AFB, DC 20332-6448	
6b. OFFICE SYMBOL (If applicable)		8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR	
8b. OFFICE SYMBOL (If applicable) NM		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620 85 C 0144	
8c. ADDRESS (City, State and ZIP Code) Building 410, Bolling AFB, DC 20332-6448		10. SOURCE OF FUNDING NOS.	
11. TITLE (Include Security Classification) Strong Representation of Weak Convergence		PROGRAM ELEMENT NO. 6.1102F	PROJECT NO. 2304
12. PERSONAL AUTHOR(S) Bai, Z.D., Liang, W.Q. and Vervaat, W.		TASK NO. A5	WORK UNIT NO.
13a. TYPE OF REPORT preprint	13b. TIME COVERED FROM 9/87 TO 9/87	14. DATE OF REPORT (Yr., Mo., Day) June 1987	15. PAGE COUNT 5
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	Keywords and phrases: Skorohod's representation theorem strong representation of weak convergence.	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
<p>The following result is proved. If <math>S_n</math> is a separable metric space for <math>n \leq \infty</math>, <math>\phi_n: S_n \rightarrow S_\infty</math> is measurable for <math>n &lt; \infty</math>, <math>X_n</math> is an <math>S_n</math>-valued random variable for <math>n \leq \infty</math> and <math>\phi_n(X_n) \rightarrow_d X_\infty</math> in <math>S_\infty</math>, then there exist <math>S_n</math>-valued random variables <math>X_n^*</math> such that <math>X_n = X_n^*</math> for <math>n &lt; \infty</math> and <math>\phi_n(X_n^*) \rightarrow_d X_\infty</math> wpl. Conditions on <math>S_n</math> and <math>\phi_n</math> are presented that allow a construction in the context of Polish spaces.</p>			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL Peegay Ravitch Maji Crowley		22b. TELEPHONE NUMBER (Include Area Code) 919-962-2307 5025	22c. OFFICE SYMBOL AFOSR/NM

AFOSR-TR. 87-1354

## CENTER FOR STOCHASTIC PROCESSES

Department of Statistics  
University of North Carolina  
Chapel Hill, North Carolina



STRONG REPRESENTATION OF WEAK CONVERGENCE

by

Z.D. Bai

W.Q. Liang

and

W. Vervaat

Technical Report No. 186

June 1987

## STRONG REPRESENTATION OF WEAK CONVERGENCE

Z.D. BAI\*

*Center for Multivariate Analysis, 515 Thackeray Hall, University of Pittsburgh, Pittsburgh, PA 15260, USA*

W.Q. LIANG\*

*Center for Multivariate Analysis, 515 Thackeray Hall, University of Pittsburgh, Pittsburgh, PA 15260, USA*

Wim VERVAAT\*\*

*Department of Mathematics, Catholic University, Toernooiveld 5, 6525 ED Nijmegen, The Netherlands*

The following result is proved. If  $S_n$  is a separable metric space for  $n \leq \infty$ ,  $q_n: S_n \rightarrow S_\infty$  is measurable for  $n < \infty$ ,  $X_n$  is an  $S_n$ -valued random variable for  $n \leq \infty$  and  $q_n(X_n) \rightarrow_d X_\infty$  in  $S_\infty$ , then there exist  $S_n$ -valued random variables  $X_n^*$  such that  $X_n^* =_d X_n$  for  $n \leq \infty$  and  $q_n(X_n^*) \rightarrow X_\infty^*$  wpl. Conditions on  $S_n$  and  $q_n$  are presented that allow a construction in the context of Polish spaces.

*AMS 1980 Subject Classification:* Primary 60B10

Skorohod's representation theorem • strong representation of weak convergence

- This research has been supported by the Air Force Office of Scientific Research Grant No. F49620-85-C-0008
- \*\* Research completed during a visit to the Center for Stochastic Processes, University of North Carolina and supported by the Air Force Office of Scientific Research Grant No. F49620-85-C-0144

In this paper we prove the following variant of Skorohod's representation theorem for weak convergence. Equality in distribution is denoted by  $=_d$ , convergence in distribution by  $\rightarrow_d$ .

**Theorem 1.** *Let  $S_n$  for  $n = 1, 2, \dots, \infty$  be a separable metric space and let  $q_n$  for  $n = 1, 2, \dots$  be a measurable function from  $S_n$  into  $S_\infty$ . If  $X_n$  is an  $S_n$ -valued random variable for  $n = 1, 2, \dots, \infty$  and  $q_n(X_n) \rightarrow_d X_\infty$  in  $S_\infty$ , then there exist  $S_n$ -valued random variables  $X_n^*$  for  $n = 1, 2, \dots, \infty$  defined on one probability space and such that  $X_n^* =_d X_n$  in  $S_n$  for  $n = 1, 2, \dots, \infty$  and  $q_n(X_n^*) \rightarrow X_\infty^*$  wpl in  $S_\infty$ .*

When  $S_n = S$  (separable) and  $q_n = \text{id}_S$  for all  $n$ , then the above theorem specializes to Dudley's (1968) variant of Skorohod's representation theorem. In Skorohod's (1956) original version  $S$  was required to be complete as well. See Wichura (1970) and Blackwell & Dubins (1983) for further extensions. Our proof of the present theorem amounts to the construction of a special metric space  $T$  to which Dudley's theorem can be applied.

Theorem 1 turns out to be useful in many instances. It is applied in Bai (1984), Bai & Yin (1986) and Yin (1984). The need for these applications led the first two authors to the present research.

Here is the simplest example of a theorem that can be proved by Theorem 1, but not by the theorem of Skorohod-Dudley in its original form. It is Theorem 4.1 of Billingsley (1968), restricted to separable metric spaces.

**Theorem 2.** *If  $S$  is a separable metric space with metric  $\rho$ ,  $(X_n, Y_n)$  are  $S^2$ -valued random variables for  $n = 1, 2, \dots$  and  $X$  is an  $S$ -valued random variable such that  $X_n \rightarrow_d X$  in  $S$  and  $\rho(X_n, Y_n) \rightarrow_d 0$  in  $\mathbb{R}$ , then  $Y_n \rightarrow_d X$  in  $S$ .*

**Proof.** By Billingsley (1968, Th.4.4) we have  $(X_n, \rho(X_n, Y_n)) \rightarrow_d (X, 0)$  in  $S \times \mathbb{R}$ . Apply Theorem 1 with  $S = S \times \mathbb{R}$ ,  $S_n = S^2$ ,  $X_n$  replaced by  $(X_n, Y_n)$  and  $q_n(x, y) = (x, \rho(x, y))$ , all for  $n < \infty$ .  $\square$

**Proof of Theorem 1.** All statements involving  $n$  are supposed to hold for  $n = 1, 2, \dots, \infty$  unless restricted explicitly; limit statements without explicit tendency hold as  $n \rightarrow \infty$ . Let  $T$  be the disjoint union of all  $S_n$ . Let  $s: T \rightarrow \{1, 2, \dots, \infty\}$  be defined by  $s(x) = n$  if  $x \in S_n$ . Set  $q_n := \text{id}_S$  and define  $q: T \rightarrow S$  by  $q(x) := q_{s(x)}(x)$ . Let  $\rho_n$  be the metric of  $S_n$ . Let  $\varepsilon_n$  be positive for  $n < \infty$ , decreasing to 0 as  $n \rightarrow \infty$ , and set  $\varepsilon_\infty := 0$ . We now define what is going to be the metric on  $T$ :

$$(1) \quad \delta(x, y) := \rho_{s(x)}(q_{s(x)}(x), q_{s(y)}(y)) + \begin{cases} \varepsilon_{s(x)} \wedge \rho_{s(x)}(x, y) & \text{if } s(x) = s(y), \\ \varepsilon_{s(x)} \vee \varepsilon_{s(y)} & \text{if } s(x) \neq s(y). \end{cases}$$

Let us first verify that  $\delta$  is indeed a metric. Obviously,  $\delta(x, y) = \delta(y, x)$  and  $\delta(x, x) = 0$ . If  $\delta(x, y) = 0$ , then  $s(x) = s(y)$  and  $\rho_{s(x)}(x, y) = 0$ , so  $x = y$ . The triangle inequality can be verified separately for both terms on the right-hand side of (1). We note the following properties of  $\delta$ -convergence:

(2)  $\delta = \rho_n$  on  $S_n \times S_n$ .

(3) If  $x, x_k \in S_n$ , then  $\delta(x, x_k) \rightarrow 0$  as  $k \rightarrow \infty$  iff  $\rho_n(x, x_k) \rightarrow 0$  and  $\rho_n(q_n(x), q_n(x_k)) \rightarrow 0$  as  $k \rightarrow \infty$ . It follows that the  $\rho_n$ -topology in  $S_n$  is coarser than or equal to the trace in  $S_n$  of the  $\delta$ -topology in  $T$ , which is homeomorphic via  $S_n \ni x \mapsto (x, q_n(x))$  to the trace in the graph of  $q_n$  of the product topology in  $S_n \times S_n$ . The last topology is separable, and so is the trace of the  $\delta$ -topology in each  $S_n$ . Consequently,  $\delta$  is separable.

(4)  $S_n$  is  $\delta$ -open for  $n < \infty$ .

(5) If  $x_n \in S_n$  for each  $n < \infty$ , then  $x_n \rightarrow x$  iff  $x \in S_\infty$  and  $q_\infty(x_n) \rightarrow x$  in  $S_\infty$ .

Having established that  $T$  with  $\delta$  is a separable metric space, we may apply Dudley's theorem to  $T$ -valued random variables. However, there is one more barrier to take. We want to identify  $S_n$ -valued random variables with  $T$ -valued random variables having range in  $S_n$ . In the first appearance random variables must be  $\mathcal{S}_n$ -measurable, where  $\mathcal{S}_n$  is the Borel field in  $S_n$  generated by  $\rho_n$ . In the

second appearance they must be  $\tilde{\sigma}_n$ -measurable, where  $\tilde{\sigma}_n$  is the trace in  $S_n$  of  $\tilde{\sigma}$ , the Borel field in  $T$  generated by  $\delta$ . So we must prove  $S_n = \tilde{\sigma}_n$ .

From the second clause in (3) it follows that  $S_n \subset \tilde{\sigma}_n$ . For the converse inclusion we must do a little more. First note that  $\tilde{\sigma}$  is already generated by the open  $\delta$ -balls in  $T$ , since  $\delta$  is separable. This can be phrased equivalently by stating that  $\tilde{\sigma}$  is the smallest  $\sigma$ -field in  $T$  which makes the functions  $\delta(x, \cdot)$  measurable for all  $x \in T$ . Consequently,  $\tilde{\sigma}_n$  is the smallest  $\sigma$ -field in  $S_n$  which makes the functions  $\delta(x, \cdot)$ , restricted to  $S_n$ , measurable for all  $x \in T$ . First suppose  $n = \infty$ . Then  $\delta(x, y) = \varrho_n(q(x), y) + \varepsilon_{n+1}$  for  $x \in T, y \in S_n$ , which, as a function of  $y$ , is obviously  $S_n$ -measurable. So  $\tilde{\sigma}_n \subset S_n$ . Considering (1) for  $x \in T$  as a function of  $y \in S_n$  we observe that  $\varrho_n(x, \cdot)$  is  $S_n$ -measurable, and that  $\varrho_n(q(x), q_n(\cdot))$  is  $S_n$ -measurable as composition of the  $S_n$ - $S_n$ -measurable function  $q_n$  and the  $S_n$ -measurable function  $\varrho_n(q(x), \cdot)$ .

We now write down the scheme of implications that proves the theorem. We are given  $S_n$ -valued random variables  $X_n$  such that

$$(6) \quad q_n(X_n) \rightarrow_d X \quad \text{in } S_n.$$

The major point, to be proved below, is that this implies

$$(7) \quad X_n \rightarrow_d X \quad \text{in } T.$$

By Dudley's theorem there are  $T$ -valued random variables  $Y_n$ , defined on one probability space, such that  $Y_n =_d X_n$  in  $T$  and  $Y_n \rightarrow Y$  wpl in  $T$ . By (4) we have  $S_n \in \tilde{\sigma}$  for each  $n$ , so there is a measurable function  $j_n: T \rightarrow S_n$  such that the restriction of  $j_n$  to  $S_n$  is the identity map on  $S_n$ ; take  $j_n$  to be the identity map on  $S_n$  and constant on  $T \setminus S_n$ . Set  $X_n^* := j_n Y_n$ . Then  $X_n^*$  has range in  $S_n$ , and  $X_n^* = Y_n$  wpl in  $T$ , since  $P[Y_n \in S_n] = P[X_n \in S_n] = 1$ . From  $X_n^* = Y_n$  wpl and  $Y_n =_d X_n$  in  $T$  it follows that  $X_n^* =_d X_n$  in  $T$ . As  $X_n^*$  has range in  $S_n$ , this implies  $X_n^* =_d X_n$  in  $S_n$ . From  $X_n^* = Y_n$  wpl in  $T$  and  $Y_n \rightarrow Y$  wpl in  $T$  we obtain  $X_n^* \rightarrow X^*$  wpl in  $T$ . As  $X_n^*$  has range in  $S_n$ , this implies by (5)

$$q(X_n^*) \rightarrow X^* \text{ wpl in } S_n.$$

We have arrived at all conclusions of the theorem.

It remains to prove the implication (6)  $\Rightarrow$  (7). We will interpret (6) and (7) by convergence of probability distributions on continuity sets, so we must compare the boundaries under  $\varrho_n$  and  $\delta$ . By (2) we have for  $A \subseteq T$ :

$$(8) \quad \partial_n(A \cap S_n) = \partial_{S_n}(A \cap S_n) = \partial_S(A).$$

Let  $B(x, \varepsilon) := \{y \in T: \delta(x, y) < \varepsilon\}$  and set

$$(9) \quad \mathfrak{V} := \{B(x, \varepsilon): \varepsilon < \varepsilon_{n+1} \text{ if } s(x) < \infty, P[X_n \in \partial_S B(x, \varepsilon)] = 0\},$$

and let  $\mathcal{U}$  consist of the unions of finitely many elements of  $\mathfrak{V}$ . By Billingsley (1968, Corollary 2 on p 15) it is sufficient for (7) that

$$(10) \quad P[X_n \in A] \rightarrow P[X \in A] \quad \text{for } A \in \mathcal{U}.$$

If  $B(x, \varepsilon) \in \mathfrak{V}$  with  $s(x) < \infty$ , then  $B(x, \varepsilon) \subset S_{n+1}$ , so  $P[X_n \in B(x, \varepsilon)] = 0$  unless  $n = s(x)$ . If  $B(x, \varepsilon) \in \mathfrak{V}$  with  $x \in S_n$ , then

$$[X_n \in B(x, \varepsilon)] = [q_n(x, q_n(X_n)) < \varepsilon - \varepsilon_n] = [q_n(X_n) \in B(x, \varepsilon - \varepsilon_n) \cap S_\varepsilon],$$

so

$$(11) \quad [q_n(X_n) \in B(x, \varepsilon) \cap S_\varepsilon] \subset \liminf [X_n \in B(x, \varepsilon)] \subset \limsup [X_n \in B(x, \varepsilon)] \\ \subset [q_n(X_n) \in \overline{B(x, \varepsilon) \cap S_\varepsilon}].$$

By (6), (8) and (9) the outmost sides of (11) have equal probabilities. Combining the previous observations for separate  $B(x, \varepsilon) \in \mathfrak{V}$  we arrive at (11) with  $A \in \mathcal{U}$  instead of  $B(x, \varepsilon) \in \mathfrak{V}$ , again with equal probabilities for the outmost sides. This proves (10), hence (7). The proof of the theorem is complete.

**Remarks.** In general the space  $T$  is not complete under  $\delta$ , even if all  $S_n$  are under  $q_n$ . To see this, consider the case that all  $x_n$  ( $n < \infty$ ) lie in  $S_m$  for one fixed  $m$ . Then  $(x_n)$  is  $\delta$ -Cauchy iff  $((x_n, q_n(x_n)))_{n < \infty}$  is  $q_m \times q_m$ -Cauchy. If the latter holds, then  $((x_n, q_m(x_n)))$  converges in  $S_m \times S_\varepsilon$ , but not necessarily in  $\text{graph } q_m$ , unless the latter is closed. This combined with the observation that  $\delta$ -Cauchy sequences  $(x_n)_{n < \infty}$  with  $x_n \in S_n$  converge if  $S_\varepsilon$  is  $q_\varepsilon$ -complete leads us to the following result.

**Theorem 3.** *Let  $S_n$  be separable and  $q_n$ -complete for each  $n$ . Then  $T$  is  $\delta$ -complete iff  $\text{graph } q_n$  is closed in  $S_n \times S_\varepsilon$  for each  $n$ .*

It is well-known that  $\text{graph } q_n$  is closed if  $q_n$  is continuous, and that  $q_n$  is continuous if  $\text{graph } q_n$  is closed and  $S_\varepsilon$  is compact. Using the fact that a subset of a Polish space is Polish iff it is  $G_\delta$  (Dugundji (1966, Th.XIV.8.3)), we arrive at the following variation on Theorem 3.

**Theorem 4.** *Let  $S_n$  be Polish for each  $n$ . Then  $T$  is Polish iff  $\text{graph } q_n$  is  $G_\delta$  in  $S_n \times S_\varepsilon$  for each  $n$ .*

For results on real functions with  $G_\delta$  graphs, see van Rooij & Schikhof (1982, Exerc. 11.Y,Z). Functions of the first class of Baire (pointwise limits of continuous functions) have  $G_\delta$  graphs.  $F_\sigma$  graphs are also  $G_\delta$ .

**Acknowledgment.** Thomas M. Liggett has contributed to our discussions about this proof.



## References

- [1] BAI, Z. D. (1984) A note on the joint distribution of the eigenvalues of noncentral multivariate  $F$ -matrix. Center for Multivariate Analysis, University of Pittsburgh. Technical Report 84-49.
- [2] BAI, Z. D., & Y. Q. YIN (1986). Limiting behavior of the norm of products of random matrices and two problems of Geman and Hwang. *Probab. Theory Rel. Fields* **73** 555-570.
- [3] BILLINGSLEY, P. (1968). *Convergence of Probability Measures*. Wiley.
- [4] BLACKWELL, D. & I. E. DUBINS (1983). An extension of Skorohod's almost sure representation theorem. *Proc. Amer. Math. Soc.* **89** 691-692.
- [5] DUDLEY, R. M. (1968). Distances of probability measures and random variables. *Ann. Math. Statist.* **39** 1563-1572.
- [6] DUGUNDJI, J. (1966). *Topology*. Allyn and Bacon.
- [7] SKOROHOD, A. V. (1956). Limit theorems for stochastic processes. *Theory Probab. Appl.* **1.3** 261-290.
- [8] VAN ROOIJ, A. C. M., & W. H. SCHIKHOF (1982). *A Second Course on Real Functions*. Cambridge University Press.
- [9] WICHURA, M. J. (1970). On the construction of almost uniformly convergent random variables with given weakly convergent image laws. *Ann. Math. Statist.* **41** 284-291.
- [10] YIN, Y. Q. (1984). Limiting spectral distributions for a class of random matrices. Center for Multivariate Analysis, University of Pittsburgh. Technical Report 84-50.



SEARCHED		SERIALIZED		INDEXED		FILED	
MAR 1984		MAR 1984		MAR 1984		MAR 1984	
FBI - NEW YORK		FBI - NEW YORK		FBI - NEW YORK		FBI - NEW YORK	
A-1							

END

DATE

FILMED

DEC.

1987